

Dr. Duke's Phytochemical and Ethnobotanical Database

List of Plants for (Z)-BETA-OCIMENE

Plant	Part	Low PPM	High PPM	StdDev	Reference
<i>Agastache foeniculum</i>	Plant	3	120	1.4074844462950056	--
<i>Alpinia galanga</i>	Leaf	--	1705	1.4142018319896454	--
<i>Alpinia galanga</i>	Rhizome	8	985		--
<i>Apium graveolens</i>	Seed	6	9		Charalambous, G. (Ed.). 1994. Spices, Herbs and Edible Fungi. Elsevier Science B. V. Amsterdam. 764 pp.
<i>Calamintha nepeta</i>	Shoot	9	9	-0.40626642590156903	Kirimer, N., Baser, K.H.C., Ozek, T. and Kurkcuoglu, M. 1992. Composition of the Essential Oil of Calamintha nepeta subsp. glandulosa. J. Ess. Oil Res. 4:189-190
<i>Carica papaya</i>	Fruit	--	--		--
<i>Carum carvi</i>	Seed	--	--		--
<i>Cinnamomum verum</i>	Leaf	--	0	-0.7120892822452273	Mallavarupu, G. R. et al. 1995. Investigation of the essential oil of cinnamon leaf grown at Bangalore and Hyderabad. Flav. & Fragr. J., 10: 239-242.
<i>Citrus limon</i>	Petiole	--	3.5		--
<i>Citrus paradisi</i>	Pericarp	--	1		--
<i>Citrus reticulata</i>	Fruit	--	0.1	-0.7485469887599769	--
<i>Citrus sinensis</i>	Fruit	2	3	-0.6648405774204937	--
<i>Dictamnus albus</i>	Shoot	15	15	-0.37869400950362786	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2
<i>Dictamnus albus</i>	Shoot	--	0.5	-0.4453273491319857	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Planta Med. 60:481-2

Plant	Part	Low PPM	High PPM	StdDev	Reference
Dictamnus albus	Shoot	--	15	-0.37869400950362786	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of <i>Dictamnus albus</i> from Turkey. <i>Planta Med.</i> 60:481-2
Elsholtzia blanda	Shoot	18	18	-0.3649078013046573	Bestman, H.J., Rauscher, J., Vostrowsky O., Pant, A.K., Dev. V., Perihar, R. and Mathela, C.S. 1992. Constituents of the Essential Oil of <i>Elsholtsia blanda</i> Benth. (Labiatae). <i>J. Ess. Oils Res.</i> 4: 121-124
Hyptis suaveolens	Shoot	3	3	-0.4338388422995102	Mallavarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of <i>Hyptis suaveolens</i> (L.) Poit. <i>J. Ess. Oil Res.</i> 5: 321.
Hyptis suaveolens	Shoot	--	3	-0.4338388422995102	Mallavarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of <i>Hyptis suaveolens</i> (L.) Poit. <i>J. Ess. Oil Res.</i> 5: 321.
Hyssopus officinalis	Shoot	690	690	2.7232028352647544	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two <i>Hyssopus officinalis</i> cultivars. <i>J. Ess. Oil Res.</i> 5: 609-611.
Hyssopus officinalis	Shoot	--	980	4.055869627831911	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two <i>Hyssopus officinalis</i> cultivars. <i>J. Ess. Oil Res.</i> 5: 609-611.
Leonotis leonurus	Se	24	24		Pedro, L.G., Barroso, J.G., Marques, N.T., Ascensao, L., Pais, M.S.S. and Scheffer, J.J.C. 1991. Composition of the Essential Oil from Sepals of <i>Leonotis leonurus</i> R. Br. <i>J. Ess. Oil Res.</i> 3: 451-3
Lindera benzoin	Fruit	55	75	1.413387566180471	--
Minthostachys mollis	Shoot	4	8	-0.4108618286345592	Alkire, B.H., Tucker, A.O., and Maciarello, M.J. 1994. Tipo (<i>Minthostachys mollis</i> (Lamiaceae): An Ecuadorian Mint. <i>Econ. Bot.</i> 48(1): 60-64.
Nepeta racemosa	Shoot	33	33	-0.29597676030980435	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of <i>Nepeta racemosa</i> Lam. <i>J. Ess. Oil Res.</i> 5: 215-7.
Nepeta racemosa	Shoot	--	33	-0.29597676030980435	Baser, K.H.C., Ozek, T., Akgul, A. and Tumen, G. 1993. Composition of the Essential Oil of <i>Nepeta racemosa</i> Lam. <i>J. Ess. Oil Res.</i> 5: 215-7.
Origanum sipyleum	Shoot	205	205	0.49443250976450925	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. <i>J. Ess. Oil Res.</i> 4: 139-142.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Origanum sipyleum	Shoot	--	15	-0.37869400950362786	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. <i>J. Ess. Oil Res.</i> 4: 139-142.
Origanum sipyleum	Shoot	--	205	0.49443250976450925	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. <i>J. Ess. Oil Res.</i> 4: 139-142.
Origanum sipyleum	Shoot	--	7	-0.4154572313675494	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. <i>J. Ess. Oil Res.</i> 4: 139-142.
Origanum vulgare	Plant	20	20	-0.4988150294995841	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
Origanum vulgare	Plant	7	7	-0.7466339613528808	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
Origanum vulgare	Plant	5	5	-0.7847599508687727	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
Origanum vulgare	Plant	5	5	-0.7847599508687727	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four <i>Origanum vulgare</i> Subspecies of Anatolian Origin. <i>J. Ess. Oil Res.</i> , 5: 425-431.
Pelargonium citrosum	Shoot	--	3	-0.4338388422995102	Matsuda, B. M., et al. 1996. Essential Oil Analysis and Field Evaluation of the Citrosa Plant ' <i>Pelargonium citrosum</i> ' as a Repellent Against Populations of <i>Aedes</i> Mosquitoes. <i>J. Am. Mosq. Contr. Assoc.</i> 12(1):69-74.
Porophyllum ruderale	Plant	80	120	1.4074844462950056	Loayza, I., de Groot, W., Lorenzo, D. et al. 1999. Composition of the essential oil of <i>Porophyllum ruderale</i> (Jacq.) Cass. from Bolivia. <i>Flav. & Fragr. J.</i> 14: 393-8.
Rosmarinus eriocalyx	Shoot	0.1	0.1	-0.4471655102251818	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the <i>Rosmarinus eriocalyx-tomentosus</i> Complex in Southeast Spain. <i>J. Ess. Oil Res.</i> 5: 243-246.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Rosmarinus eriocalyx	Shoot	--	--		Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	6	75	-0.10296984552421623	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	--	--		Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	--	13	-0.38788481496960825	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	6	75	-0.10296984552421623	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus tomentosus	Shoot	0.1	0.1	-0.4471655102251818	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus x lavandulaceus	Shoot	26	110	0.05786925013044062	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus x mendizabalii	Shoot	0.1	0.1	-0.4471655102251818	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Sassafras albidum	Leaf	2.6	8	-0.7021125497444185	--
Satureja cilicica	Shoot	210	210	0.5174095234294602	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of Satureja cilicica P.H. Davis. J. Ess. Oil Res. 5: 547-548.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Tagetes lucida	Shoot	--	--		Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of Tagetes lucida Cav. ssp. lucida essential oil. Flavor & Fragrance, 12(1): 47-52.
Thymus longicaulis	Shoot	0.8	0.8	-0.4439487283120886	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	0	-0.4476250504984808	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.
Thymus longicaulis	Shoot	--	0.8	-0.4439487283120886	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of Thymus longicaulis C. Presl subsp. longicaulis in the same Population. J. Ess. Oil Res. 5: 291-5.